


AN 91-356487 [49] WPIDS
 DNN N91-272859
 TI **Brush** assembly for fractional horsepower DC **motor**
 - has two **brushes** side by side connected electrically in
 parallel and having different **resonant frequencies**
 .
 DC V04 V06
 IN BALNES, R F; BAINES, R F
 PA (JOHN-N) JOHNSON ELECTRIC SA
 CYC 3
 PI GB 2244603 A 911204 (9149)*
 JP 05146108 A 930611 (9328) H02K005-14
 GB 2244603 B 940727 (9427) 2 pp H02K013-00
 ADT GB 2244603 A GB 90-11275 900521; JP 05146108 A JP 91-114938 910520;
 GB 2244603 B GB 90-11275 900521
 PRAI GB 90-11275 900521
 IC H01R039-62; H02K013-00
 ICM H02K005-14
 ICS H01R039-62; H02K013-00; H02K023-00
 / BINARY DATA / IMAGE001.TIF
 AB GB 2244603 A UPAB: 930928
 The electric **motor brush** assembly comprises
 resilient electrically conductive support arranged to carry two or
 more **brushes** axially displaced with respect to a
 longitudinal axis of the **motor** and connected electrically
 in parallel. The support normally comprises a separate arm (18,19)
 for each **brush** ((20,21)).
 The separate arms may be arranged to have different natural
 resonance frequencies of oscillation. The **brushes** may be
 different sizes and/or of different physical densities.
 ADVANTAGE - Reduces current density required for each
brush without increasing size of **brushes**.
 3/6
 FS EPI
 FA AB; GI
 MC EPI: V04-L01B; V06-M12

CRC Handbook of Chemistry and Physics

A Ready-Reference Book of Chemical and Physical Data



Editor

Robert C. Weast, Ph.D.

Formerly Vice President Research, Consolidated Natural Gas Service Company, Inc.
Formerly Professor of Chemistry at Case Institute of Technology

Associate Editor

Melvin J. Astle, Ph.D.

Formerly Professor of Organic Chemistry at Case Institute of Technology
and
Manager of Research at Glidden-Durkee Division of SCM Corporation

In collaboration with a large number of professional chemists and physicists whose assistance is acknowledged in the list of general collaborators and in connection with the particular tables or sections involved.



CRC Press, Inc.
Boca Raton, Florida

© 1981

PHYSICAL CONSTANTS OF INORGANIC COMPOUNDS (Continued)

Other solvents	Name	Synonyms and Formulae	Mol. wt.	Crystalline form, properties and index of refraction	Density or spec. gravity	Melting point, °C	Boiling point, °C	Solubility, in grams per 100 cc		
								Cold water	Hot water	Other solvents
	Cobalt complexes									
	Ethylenediaminecobalt(III) chloride	$\text{Co}(\text{C}_2\text{H}_4(\text{NH}_2)_2)_3\text{Cl}_3 \cdot 3\text{H}_2\text{O}$	399.64	br pr.	1.542 ¹⁷	256; -3H ₂ O, 100		v s		
	Triethylenediaminecobalt(III) chloride	$\text{Co}(\text{NH}_2)_3(\text{NO}_2)_3$	248.04	yel, rhomb pl or leaf	1.992 ¹⁸	d 158	exp 164	0.177 ^{18,19}	0.28 ²⁰	
	Triethylenediaminecobalt(III) nitrate	$[\text{Co}(\text{NH}_2)_3(\text{NO}_2)_3]\text{NO}_3$	265.07	yel, rhomb	1.922 ¹⁷			3 ²⁰		
	Potassium tetrakisethylenediaminecobaltate(III)	$\text{K}[\text{Co}(\text{NH}_2)_4(\text{NO}_2)_4]$	316.12	yel, rhomb	2.076 ¹⁸			1.758 ^{18,19}		
	Columbium	see Niobium.	68.546	redsh met, cub.	8.92	1063.4±0.2	2567	i	i	s HNO ₃ , h H ₂ SO ₄ ; v s a; HCl, NH ₄ OH; dil a, NH ₄ OH; al s; 7.14 al; s eth
	Copper									
	Blue verdigris.	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot \text{CuO} \cdot 0.5\text{H}_2\text{O}$	369.26	grnsh-bl powd.				sl s		
	Neutral verdigris.	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot \text{H}_2\text{O}$	199.65	dk grn powd, 1.545, 1.550	1.882, anhydr-1.93	115	d 240	7.2	20	
	Paris green.	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 3\text{Cu}(\text{AsO}_2)_2$ (approx)	1013.77	em grn powd.				i		s a, NH ₄ OH; i al
	(II) acetate	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2$	151.10	red, amorph, expl.		exp d 100-105	exp 202	v s	d	s a, KCN; d a; i MeOH
	(II) acetylacetonate	$\text{Cu}(\text{NH}_2)_2(\text{N}_3)_2$	181.64	dk grn cr, exp		260-270	d 300	i		s NH ₄ OH; i abs a
	(II) diamminechloride, di-	$\text{Cu}(\text{NH}_3)_2\text{Cl}_2$	168.51	grn cr.	2.32 ¹⁸			v s		
	(II) diamminechloride, di-	$\text{Cu}(\text{NH}_3)_2\text{Cl}_2$	236.63	bl, cub.	1.48 ¹⁸			s	d	
	(II) diamminechloride, di-	$[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$	291.79	vit-bl cr.		d 160		s		
	(II) diamminechloride, di-	$[\text{Cu}(\text{NH}_3)_4](\text{NO}_3)_2$	255.67	dk-bl, oct.	1.91 ¹⁸	d 210 exp		s		
	(II) diamminechloride, di-	$[\text{Cu}(\text{NH}_3)_4](\text{NO}_3)_2$	223.61	vit-bl, tetr.		-2NH ₃ , 97		v s		
	(II) diamminechloride, di-	Cuprum ammoniacale. $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4 \cdot \text{H}_2\text{O}$	245.74	dk-bl, rhomb, unstab	1.79 ¹⁸	-NH ₃ ·H ₂ O, 30		18.05 ^{18,19}		
	(II) diamminechloride, di-	CuSb	312.37	gray	8.51	687		i	i	s a, NH ₄ OH
	(II) diamminechloride, di-	$\text{Cu}_3(\text{AsO}_4)_2 \cdot 4\text{H}_2\text{O}$	540.52	blsh-grn				i		s a, NH ₄ OH
	(II) diamminechloride, di-	$\text{CuH}_2(\text{AsO}_4)_2 \cdot 2\text{H}_2\text{O}$	911.42	bl				i		s a, NH ₄ OH
	(II) diamminechloride, di-	CuAs	467.54	bl, oct.	7.56			i		s a, NH ₄ OH; i al
	(II) diamminechloride, di-	Nat. domoykite. CuAs	265.54	hex	8.0	830		i		s a, NH ₄ OH; i al
	(II) diamminechloride, di-	Scheele's green. $\text{CuHAsO}_3(?)$	187.47	grn powd.		d				d conc H ₂ SO ₄ ; s NH ₄ Cl; v s dil a
	(II) diamminechloride, di-	CuN_3	105.56	col cr, v exp	3.26			0.00075 ²⁰		
	(II) diamminechloride, di-	$\text{Cu}(\text{N}_3)_2$	147.58	brn-red or brn-yel cr, exp	2.604	exp 215		0.008 ²⁰		s dil a; al s al
	(II) diamminechloride, di-	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$	341.80	lt bl cr powd.		-H ₂ O, 110		sl s		
	(II) diamminechloride, di-	$\text{Cu}(\text{BO}_2)_2$	149.16	blsh grn cr powd.	3.859					s NH ₄ OH
	(II) diamminechloride, di-	$\text{Cu}_2\text{B}_2\text{O}_7$	212.24	yel	8.116					s HBr, HCl, HNO ₃ , NH ₄ OH; i acet
	(II) diamminechloride, di-	$\text{Cu}(\text{BrO}_3)_2 \cdot 6\text{H}_2\text{O}$	427.45	bl-grn, cub.	2.583	d 180	-6H ₂ O, 200	v s	d	s al, acet, NH ₃ , pyr; i ba
	(II) diamminechloride, di-	CuBr (or Cu_2Br_2)	143.45	wh, cub, 2.116	4.98	492	1345	v s		s dil min a, NH ₄ OH; v s a; s al, eth, NH ₄ OH, dil a
	(II) diamminechloride, di-	CuBr_2	223.31	blk, monoc, deliq	4.77 ¹⁸	498		v s		s a, NH ₄ OH; 0.026 aq CO ₂ ; s a, NH ₄ OH, KCN; i al
	(II) diamminechloride, di-	$\text{CuBr}_2 \cdot 3\text{Cu}(\text{OH})_2$	516.02	em grn, rhomb	4.00	-H ₂ O, 210-d 240-250		i	d	s NH ₄ OH, h NaHCO ₃
	(II) diamminechloride, di-	$\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$	273.77	dk grn cr.				v s		s al, acet
	(II) diamminechloride, di-	Cu_2CO_3	187.09	yel	4.40	d 200		i		s dil a
	(II) diamminechloride, di-	Nat. malachite. $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	221.11	dk grn, monoc, 1.655, 1.875, 1.909	4.0			i		
	(II) diamminechloride, di-	Nat. azurite, chrysylite. $2\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	344.65	bl, monoc, 1.730, 1.758, 1.838	3.88	d 220		i		
	(II) diamminechloride, di-	$\text{Cu}(\text{ClO}_3)_2 \cdot 6\text{H}_2\text{O}$	338.53	grn, cub, deliq		65	d 100	207 ²⁰	v s	
	(II) diamminechloride, di-	$\text{Cu}(\text{ClO}_3)_2 \cdot 3\text{Cu}(\text{OH})_2$	523.11	grn cr or amorph.	3.55	d		i		

PHYSICAL CONSTANTS OF INORGANIC COMPOUNDS (Continued)

	Name	Synonyms and Formulae	Mol. wt.	Crystalline form, properties and index of refraction	Density or spec. gravity	Melting point, °C	Boiling point, °C	Solubility, in grams per 100 cc		
								Cold water	Hot water	Other solvents
00 cc	Calcium	$\text{Ca}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$	350.34	wh, oct.		-2H ₂ O, 120		4 ²⁵	s	s al
solvents	salicylate	CaSeO_4	183.04	col.	2.88			7.9 ⁴	5.4 ⁶⁷	
oet	selenate	$\text{CaSeO}_4 \cdot 2\text{H}_2\text{O}$	219.07	col, monoc.	2.68					
	selenate, dihydrate	$\text{CaSeO}_4 \cdot 2\text{H}_2\text{O}$	119.04	cub, 2.274	3.57			0.0095 ¹⁷		s HCl
	selenide	CaSe	116.16	col, monoc, 1.610	2.905	1540				
	metasilicate (α)	Nat. pseudowollastonite. CaSiO_3		1.611, 1.664						
	metasilicate (β)	Nat. wollastonite. CaSiO_3	116.16	col, monoc, 1.616	2.5	tr 1200				
al, eth	di-orthosilicate (I)	Ca_2SiO_4	172.24	col, monoc, 1.717	3.27	2130				
1.72 ²⁵ al	di-orthosilicate (II)	Ca_2SiO_4	172.24	col, rhomb, 1.717	3.28	tr to (I) 1420				
	di-orthosilicate (III)	Ca_2SiO_4	172.24	col, monoc, 1.642	2.97	tr to 675				
	(tri-)silicate	Nat. alite. Ca_3SiO_5 or $(3\text{CaO} \cdot \text{SiO}_2)$	228.32	col, monoc, α 1.718, β 1.724		1900 (incogr)				s a, alk
	silicide	CaSi	96.25	cr powd.	2.5	179-180		0.004 ¹⁵	d	i al, eth
	stearate	$\text{Ca}(\text{C}_{18}\text{H}_{35}\text{O}_2)_2$	607.04	col, rhomb, 1.569				0.193 ¹⁰	0.89 ²⁰	
	succinate	$\text{CaC}_4\text{H}_4\text{O}_6 \cdot 3\text{H}_2\text{O}$	212.22	col, rhomb, 1.575, 1.613				0.209 ²⁰	0.1619 ¹⁰⁰	s a, NH ₄ salts, Na ₂ SO ₄ , glyc
	sulfate	Nat. anhydrite. CaSO_4	136.14	col, hex or tric, 1.505, 1.548	2.61	tr to rhomb >200				
	sulfate	Soluble anhydrite. CaSO_4	136.14	wh powd.		-1/2 H ₂ O, 163		0.3 ²⁰	al s	s a, NH ₄ salts, Na ₂ SO ₄ , glyc
	sulfate half-hydrate	Plaster of Paris. $\text{CaSO}_4 \cdot 1/2\text{H}_2\text{O}$	145.15					0.241	0.222 ¹⁰⁰	s a, NH ₄ salts, Na ₂ SO ₄ , glyc
OH	sulfate dihydrate	Nat. gypsum. $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	172.17	col, monoc, 1.521	2.32	-1 1/2 H ₂ O, 128		0.021 ¹⁴ d	0.048 ²⁰ d	d a
al, eth	sulfide	Nat. oldhamite. CaS	72.14	col, cub, 2.137	2.5	d 15-18		0.0043 ¹⁵	0.0011 ¹⁰⁰	s al, s H ₂ SO ₄
al: s MeOH?	sulfide, hydro-	$\text{Ca}(\text{HS})_2 \cdot 8\text{H}_2\text{O}$	214.32	col, hex		-WH ₂ O, >150				s a
NH ₄ acet?	sulfite	$\text{Ca}(\text{SO}_3) \cdot 1/2\text{H}_2\text{O}$	129.15	yellow liq, strong SO ₂ odor				0.0266 ⁹	0.0689 ^{17,18}	al s al
ch	sulfite, dihydrogen	$\text{Ca}(\text{HSO}_3)_2$	202.22	col, rhomb, 1.525, 1.535, 1.550		d		0.0032 ²	0.0078 ^{17,18}	s HCl; i ac, a
acet	d-tartrate	$\text{CaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$	260.21	tric, powd or need		-4H ₂ O, 200			0.16 ¹⁰⁰	0.28 ¹⁵ , 0.85 ¹⁰⁰ ac a
	dl-tartrate	$\text{CaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$	260.21	wh, monoc or tric pr		-3H ₂ O <170				
al; d abs al?	mesotartarate	$\text{CaC}_4\text{H}_4\text{O}_6 \cdot 3\text{H}_2\text{O}$	242.20	cub, 2.51, 2.58	4.873	>960		al s	s	s a
al	telluride	CaTe	167.68	wh fl.				s	s	i al
	tellurite	CaTeO_3	215.68	yel cr				v s	v s	v s al
	thiocarbonate, tri-	CaCS_3	148.28	wh cr, deliq				16 ⁹	30 ²⁰	s al
eth	thiocyanate	$\text{Ca}(\text{SCN})_2 \cdot 3\text{H}_2\text{O}$	210.29	col, trig, 1.5496	2.176			100 ²		
i ac a	di-thionate	$\text{Ca}(\text{SO}_3)_2 \cdot 4\text{H}_2\text{O}$	272.27	tric, 1.872		d 1975				
i ac a	thiosulfate	$\text{CaS}_2\text{O}_3 \cdot 6\text{H}_2\text{O}$	260.30	col, cub, rhomb, β 2.34	4.10			0.00064 ¹⁵	0.00012 ¹⁰⁰	
	metatitanate	Nat. perovskite. CaTiO_3	135.98	wh, tetr, 1.9263, 1.9107	6.062 ²⁰			0.2		i al, a; s NH ₄ Cl
	tungstate	CaWO_4	287.93	col or w sc, tetr, 1.918, 1.934	6.06					d a
al, NH ₄ salts?	tungstate	Nat. scheelite. CaWO_4	287.93	col, tric.		-7H ₂ O, 105				
al, eth	metatungstate	$\text{Ca}_5\text{H}_4(\text{H}_2(\text{WO}_4)_4) \cdot 27\text{H}_2\text{O}$	3500.96					8.28 ⁹	7.39 ¹⁰⁰	
al; s al; 0.006 ²⁵	valerate	$\text{Ca}(\text{C}_4\text{H}_7\text{O}_2)_2$	242.33	col, monoc.	4.78	2550				i a, alk
th	metazirconate	CaZrO_3	179.30	col, cub, 2.4173	3.51	>3550				s liq Fe; i a, alk
al	Carbon	Diamond. C	12.01	col, hex	2.25 ²⁰	subl 3652-97				i a, alk
	carbon	Graphite. C	12.01	amorph, blk	1.8-2.1	subl 3652-97				s CS ₂ ; v s al, eth
s al	carbon, amorphous	C	12.01	rhomb pr, 1.740, 1.847, 1.863	3.823	148-149 d				s al, eth, chl
HCl	(di-)bromide, hexa-	Hexabromomethane. C ₂ Br ₆	503.48	col, monoc or oct	3.42	tr to oct 48.4; m.p. 90.1		0.024 ²⁰		
al, s a	bromide, tetra-	Tetrabromomethane. CBr ₄	331.65			57.5				s al, eth, oils
	(di-)bromide, tetra-	Tetrabromethylene. C ₂ Br ₄	343.66	col, rhmb, tric or cub	2.091	subl 187				
al; s i NH ₄ Cl	(di-)chloride, hexa-	Hexachloro ethane. C ₂ Cl ₆	236.74	col liq, 1.4601	1.5867 ²⁰	-23				s al, bz, chl, eth
al	chloride, tetra-	Tetrachloromethane. CCl ₄	153.81			-22.4				s al, eth
al	(di-)chloride, tetra-	Tetrachloroethylene. C ₂ Cl ₄	165.83	col liq, eth odor, 1.5055	1.6311 ¹⁵					